

Research Article

Intestinal Permeability Among Calves with Behavioural Disturbances

Hasan ERDOĞAN¹, Songül ERDOĞAN¹

¹Aydın Adnan Menderes University, Veterinary Faculty, Internal Medicine Department, Aydın/Türkiye

ORCID:0000-0001-5141-5108
ORCID: 0000-0002-7833-5519

Corresponce

Aydın Adnan Menderes University, Veterinary Faculty, Internal Medicine Department, Aydın/Türkiye

E- mail: songultp.09@gmail.com
Doi: 10.5281/zenodo.10474137

Abstract

The connection among disturbed intestinal ecosystem, namely dysbiosis, and central nervous system have been recognized and evidence of proofed within the last 10 years. Microbiological alterations consequently give rise to elevated intestinal permeability, and thus the perception of bacterial fragment and toxins. This could easily hasten and instigate local/systemic inflammatory respond, acting on distant organs, involving the brain. Moreover the unfication of intestinal epithelial barrier exhibits a pivotal role for the microbiota–gut–brain axis. Novel discovery on zonulin, a signifcant tight junction modifier of intestinal epithelial cells, has been recognized to participate within the blood–brain barrier functioning. In the present article the present authors' addressed the possible role of zonulin in an attempt to detect the relationship among leaky gut, neurological involvement and gut-brain axis to those of calves. Serum zonulin levels were determined by use of Sandwich ELISA. Clinical signs were recorded and noticed. Comparative analysis of serum zonulin levels (ng/mL) were deemed available were showing statistical significance ($p=0,001$) among healthy ones ($24,11 \pm 2,44$) vs. calves with behavioural disturbances ($58,03 \pm 2,47$).

Keywords: Brain, calf, gut, permeability, tight junction, zonulin.

Introduction

Emerging literature exhibited a zetsful bidirectional connection among gut microbiome and the nervous system, denoted as microbiota–gut–brain axis (Cryan et al., 2019). Several conditions influence microbial echology, involving stress, nutritional status, environmental reflection, and intestinal lumen habitants affecting neural reflection (Cryan et al., 2019, Oriach et al., 2016). The multiplex interaction among microbiota and brain is uniterupted through inflammatory compounds, neuroactive bacteriological metabolites

neurotransmitters and vagus/enteric nerves (Cryan et al., 2019. microbiota–gut–brain axis exhibited an inter-relationship among altered gut microbiome diversity and neurological diseases (oriach et al., 2016, Morais et al., 2021, Pap et al., 2022). To our knowledge there has been no study evaluating the connection of neurological disorders and gut-brain axis in calves at field conditions. Therefore our aim was to analyze the interaction among calves, intestinal permeability and neurological signs, which prompted us t oto perform this study.

Material and Method

InfographicData

The present research was conducted in 4 different commercial farms involving calves, located in Aydın Municipality among Aegean Region of Turkey. A total of 30 Holstein calves (n=21 female, 9 male) out of 119 inspected, to those of which 34-67 days old ones were involved at the present study. Solely 21 healthy calves (14 female, 7 male) were enrolled.

Blood Sampling and Analytes

Blood was taken via Vena jugularis onto anticoagulated tubes once solely. Sera was suddenly separated and transferred to where its kept. Bovine zonulin ELISA Kit (MyBiosource ELISA kits), previously used (Alic Ural et al., 2021a, Alic Ural et al., 2021b, Alic Ural et al., 2021c, Alic Ural, 2022a, Alic Ural, 2022b, Alic Ural, 2022c, Alic Ural, 2023, Alic Ural and Ural, 2023) were purchased commercially. ELISA was preferred for serum zonulin levels similar to previous methodology (Alic Ural et al., 2021a, Alic Ural et al., 2021b, Alic Ural et al., 2021c, Alic Ural, 2022a, Alic Ural, 2022b, Alic Ural, 2022c, Alic Ural, 2023, Alic Ural and Ural, 2023, Ural et al., 2023), was adopted. Briefly Sandwich ELISA methodology was evident.

Statistical analysis

The mean and standard deviation values of zonulin levels for calves in the healthy group and those with behavioral disturbances were tabulated. For the comparison of the obtained data, the Mann-Whitney U test was utilized, and instances where the p-value was less than 0.05 were deemed statistically significant. The statistical analyses and graphs were conducted using the Graphpad Prism (9.5.0, USA) software.

Results

Circulating serum zonulin levels (ng/mL) were shown in table 1 for both groups. Besides boxplot analysis were shown in Fig 1.

Table 1. Circulating serum zonulin levels among healthy and behavioural disturbance calves.

Parametre	Healthy Calves	Calves with behavioural disturbances	P value
Zonulin (ng/mL)	24,11 ± 2,44	58,03 ± 2,47	0,001




Fig. 1. Boxplot analysis of serum zonulin levels for both group of calves.

Discussion

Intestinal permeability, exhibiting a pivotal role among microbiota-gut-brain axis, permits the piercing of luminal content through the circulatory system (Bischoff et al., 2014). Intestinal epithelial layer barrier functioning is maintained through ntercellular junctions, composed of tight/adherens junction and desmosome (Camillerei et al., 2019). Through existence of leaky gut”, tight junctions and their relevant constituents patricipate (Tripathi et al., 2009, Sturgeon and Fasano, 2016). In the present study fort he first time under our era and geograp- hy, we herein showed evidence of proof for leaky gut among calves with beahvioural disturbance. About 24 years ago, Fasano and his colleagues desc- ribed the finding of zonulin, modifying paracellu- lar permeability (Fasano et al., 2000,. Moreover the influence of the zonulin pathway on the modi- fication of intestinal permeability has been deter- mined by phase 2 clinical (Hoilat et al., 2022). Des- pite the fact that the vast majority of available data regarding zonulin has been assoociated to intestinal

diseases, its significant participation in exactly all organs, including brain etc., has currently been recognized (wang et al., 2000, Rittirsch, et al., 2013, Ghanadi, K.; Naghdi 2022, Yu et al., 2021, Smecuol et al., 2005, Miranda-Ribera et al., 2022). Interestingly a novel cohort study established the association of sera zonulin levels and severity of respiratory disorder among diseased calves. In the latter study calves diagnosed with bovine respiratory disease were subjected to specific Bovine Zonulin ELISA test kits. The mean zonulin levels (ng/mL) healthy calves (21.03 ± 10.57), calves without severe pneumonia (Score <5) (39.53 ± 11.90) and calves with pneumonia (Score >5) (64.69 ± 17.98) presented statistically significant alterations ($p < 0.05$). In that study zonulin, not only limited to exhibition of intestinal inflammation, but also conducted to show lung injury (lung permeability) (Alic Ural and Ural, 2023). Another study validated the intercation among fecal scores and sera zonulin concentrations as a biomarker for intestinal inflammation among calves with diarrhea. Specific Bovine Zonulin ELISA test kits revealed diarrheic calves with score 2 and score 3 expressing ser zonulin levels 41.04 ± 4.5 and 61.27 ± 4.80 , with significant changes in comparison to healthy calves 20.35 ± 2.93 ($p < 0.05$). In that study intestinal inflammation showed correlation with serum zonulin levels (Alic Ural, 2023). Moreover a prospective field study aimed to detect zonulin value as a noninvasive selected biomarker for intestinal barrier functioning and leaky gut. Circulatory zonulin concentrations (ng/mL) showed as ($\pm$ SEM), exhibited significant alterations ($p < 0.001$) between healthy and diarrheic calves (26.43 ± 3.528 vs. 57.97 ± 4.250), respectively (Alic Ural 2022a). On the other hand zonulin levels might be affected by both in seasonal/spatial distribution at different day/night. Calves were subjected to analysis for mean zonulin levels (ng/mL) at 12 pm and 08:00 am in the January group were 28.04 ± 12.49 vs. 11.55 ± 8.45 , respectively ($p = 0.012$), whereas at June group the mean zonulin levels at 12 pm and 08.00 am were 58.54 ± 19.16 vs 24.03 ± 9.9 , respectively ($p = 0.012$). The latter study showed that heat stress could be capable of influencing intestinal integrity and seasonal dissipation of zonulin concentration (Alic Ural 2022b). In the present study in comparison to aforementioned researches serum zonulin levels (ng/mL) exhibited statistical significance ($p = 0.001$) among healthy calves (24.11 ± 2.44) vs. calves with behavioural disturbances (58.03 ± 2.47). The reason for this results

should be dept in discussed in below paragraph. Given disturbed gut microgeography diversity, namely dysbiosis, has been connected with release of intestinal zonulin, altered gut permeability, and upregulation of inflammatory compounds (Veres-Székely et al., 2023). The extension of intestinal-derived microbial compounds, toxins, and inflammatory conditions, also composed of zonulin, lastly arrive distant organs, involving the central nervous system. The latter could prone to elevated blood-brain barrier permeability, inflammation of the nervous system and behavioral alterations could selectively exaggerated through microbiota diminishing (Miranda-Ribera et al., 2022). Taking into account all aforementioned data denoting that dysbiosis and the zonulin route could participate as major factor in microbiota-gut-brain axis associated disorders (Miranda-Ribera et al., 2022, Veres-Székely et al., 2023). During SARS-CoV-2 strike during 2020, denoted as pandemic proportions beacuse of its quick spread and virulence, COVID-19 patients exhibited digestive, respiratory and neurologic clinical presentation. Given neurotropism of the reponsible virus, infection of the gut, and related mechanisms involving zonulin brain receptor and probable usage of zonulin antagonists were inspected (Llorens et al., 2021). To the present authors' knowledge there has been no previous study, at least in our country, related to zonulin and behavioural disturbances among calves. The results that we obtained would be further investigated with larger sacle population of calves.

Referances

- Aliç Ural, D. (2022a). Zonulin as a Noninvasive Selected Biomarker of Gut Barrier Function Identify and Debug Calves Suffering from Diarrhea. *International Journal of Veterinary and Animal Research*, 5(3), 2022-3.
- Aliç Ural, D. (2022b). Heat Stress and Seasonal Dissipation of Circulating Zonulin Levels Among Calves in Aydın Region. *International Journal of Veterinary and Animal Research*, 5(2), 47-49.
- Aliç Ural, D. (2022c). Leaky Gut and Giardia duodenalis Infection Associated Serum Zonulin Levels Among Calves Randomized Clinical Study. *Türkiye Klinikleri Journal of Veterinary Sciences*, 13(2), 46-50.
- Aliç Ural, D., & Ural, K. (2023). Zonulin as a Preliminary Biomarker of Lung Permeability Among Diseased Calves: Cohort Study. *Egyptian Journal of Veterinary Sciences*, 54(4), 601-607.
- Aliç Ural, D., Ural, K., Erdoğan, H., Erdoğan, S.

- (2021a). Alterations in Gut Integrity Due to Heat Stress Among Dairy Cattle of Aydin City Analytical Interpretation of Zonulin Levels within Repetitive Measurements. *International Journal of Veterinary and Animal Research*, 4(3), 111-114.
- Aliç Ural, D., Erdoğan, S., Erdoğan, H., Ural, K. (2021b). Heat stress intestinal barrier disruption and calves multidisciplinary perspective field study. *Journal of advances in vetbio science and techniques*, 6(3), 265-269.
- Aliç Ural, D., (2023). Serum Zonulin Levels and Fecal Scoring as Probable Early Predictor of Intestinal Inflammation Among Calves with Diarrhea: Cohort Study. *Turkiye Klinikleri Journal of Veterinary Sciences*, 14(1), 18-21.
- Bischoff, S.C., Barbara, G., Buurman, W., Ockhuizen, T., Schulzke, J.-D., Serino, M., Tilg, H., Watson, A., Wells, J.M. (2014). Intestinal permeability—A new target for disease prevention and therapy. *BMC Gastroenterology*, 14, 189.
- Camilleri, M. (2019). Leaky gut: Mechanisms, measurement and clinical implications in humans. *Gut*, 68, 1516–1526.
- Cryan, J.F., O’Riordan, K.J., Cowan, C.S., Sandhu, K.V., Bastiaanssen, T.F., Boehme, M., Codagnone, M.G., Cusotto, S., Fulling, C., Golubeva, A.V. (2019). The microbiota-gut-brain axis. *Physiological Reviews*, 99, 1877–2013.
- Fasano, A., Not, T., Wang, W., Uzzau, S., Berti, I., Tommasini, A., Goldblum, S.E. (2000). Zonulin, a newly discovered modulator of intestinal permeability, and its expression in coeliac disease. *Lancet*, 355, 1518–1519.
- Ghanadi, K., Naghdi, N. (2022). The Role of Zonulin as a Prognostic Biomarker in Liver Diseases: A Systematic Review. *Advancements in Life Sciences*, 9, 277–283.
- Hoilat, G.J., Altowairqi, A.K., Ayas, M.F., Alhadad, N.T., Alnujaidi, R.A., Alharbi, H.A., Alyahyawi, N., Kamal, A., Alhabeeb, H., Albazee, E., et al., (2022). Larazotide acetate for treatment of celiac disease: A systematic review and meta-analysis of randomized controlled trials. *Clinics and Research in Hepatology and Gastroenterology*, 46, 101782.
- Llorens, S., Nava, E., Munoz-Lopez, M., Sanchez-Larsen, A., & Segura, T. (2021). Neurological symptoms of COVID-19: The zonulin hypothesis. *Frontiers in Immunology*, 12, 665300.
- Miranda-Ribera, A., Serena, G., Liu, J., Fasano, A., Kingsbury, M.A., Fiorentino, M.R. (2022). The Zonulin-transgenic mouse displays behavioral alterations ameliorated via depletion of the gut microbiota. *Tissue Barriers*, 10, 2000299.
- Morais, L.H., Schreiber, H.L., IV, Mazmanian, S.K. (2021). The gut microbiota-brain axis in behaviour and brain disorders. *Nature Reviews Microbiology*, 19, 241–255.
- Oriach, C.S., Robertson, R.C., Stanton, C., Cryan, J.F., Dinan, T.G. (2016). Food for thought: The role of nutrition in the microbiota-gut-brain axis. *Clinical Nutrition Experimental*, 6, 25–38.
- Pap, D., Veres-Székely, A., Szebeni, B., Vannay, Á. (2022). PARK7/DJ-1 as a Therapeutic Target in Gut-Brain Axis Diseases. *International Journal of Molecular Sciences*, 23, 6626.
- Rittirsch, D., Flierl, M.A., Day, D.E., Nadeau, B.A., Werner, C.M.L., Wanner, G.A., Simmen, H., Fasano, A., Ward, P.A. (2011). Role of zonulin as prehaptoglobin2 in acute lung injury. *FASEB Journal*, 25, 114.1.
- Rittirsch, D., Flierl, M.A., Nadeau, B.A., Day, D.E., Huber-Lang, M.S., Grailer, J.J., Zetoune, F.S., Andjelkovic, A.V., Fasano, A., Ward, P.A. (2013). Zonulin as prehaptoglobin2 regulates lung permeability and activates the complement system. *American Journal of Physiology-Cell Physiology*, 304, L863–L872.
- Smecuol, E., Sugai, E., Niveloni, S., Vázquez, H., Pedreira, S., Mazure, R., Moreno, M.L., Label, M., Mauriño, E., Fasano, A. (2005). Permeability, zonulin production, and enteropathy in dermatitis herpetiformis. *Clinical Gastroenterology and Hepatology*, 3, 335–341.
- Sturgeon, C., Fasano, A. (2016). Zonulin, a regulator of epithelial and endothelial barrier functions, and its involvement in chronic inflammatory diseases. *Tissue Barriers*, 4, e1251384.
- Tripathi, A., Lammers, K.M., Goldblum, S., Shea-Donohue, T., Netzel-Arnett, S., Buzza, M.S., Antalis, T.M., Vogel, S.N., Zhao, A., Yang, S. (2009). Identification of human zonulin, a physiological modulator of tight junctions, as prehaptoglobin-2. *Proceedings of the National Academy of Sciences. USA*, 106, 16799–16804.
- Ural, K., Erdoğan, H., Aliç Ural, D., Paşa, S., Erdoğan, S., Ayan, A., Gültekin, M. (2023). “Enema Treatment with the Ayurvedic Herb Curcumin in Calves Infected with *Giardia duodenalis* Cohort Study. *Turkiye Klinikleri Journal of Veterinary Sciences*, 10(2), 35-45.
- Veres-Székely, A., Szász, C., Pap, D., Szebeni, B., Bokrossy, P., Vannay, Á. (2023). Zonulin as a Potential Therapeutic Target in Microbiota-Gut-Brain Axis Disorders: Encouraging Results and Emerging Questions. *International Journal of Molecular Sciences*, 24, 7548.

Wang, W., Uzzau, S., Goldblum, S.E., Fasano, A. (2000). Human zonulin, a potential modulator of intestinal tight junctions. *Journal of Cell Science.*, 113, 4435–4440.

Yu, J., Shen, Y., Zhou, N. (2021). Advances in the role and mechanism of zonulin pathway in kidney diseases. *International Urology and Nephrology*, 53, 2081–2088.

Supplements

Table 2. Table involved calves enrolled at the present study. Clinical signs were revised and adopted from well described data (https://www.michigan.gov/mdard/-/media/Project/Websites/mdard/documents/animals/exhibitions/signs/illness_barn_posters.pdf?rev=f4f10dec01eb4a60bd7a9613131e40c9&hash=D70E252629173B4A72B403184EB4F28D)

Neurological Signs:	
N=2	Behavioral alterations or easily startled
N=3	Restlessness/agitation
N=9	Lack of coordination or high stepping
N=12	Head rubbing, tossing or pressing
N=4	Exaggerated blinking and chewing movements
N=6	Trembling or convulsions